

Flavor independent systematics of excited baryons and intra-band transition*

M. Koma (Takayama),^{a,b} A. Hosaka^b and H. Toki^b

^a Max-Planck-Institut für Physik, Föhringer Ring 6, D-80805 München, Germany

^bResearch Center for Nuclear Physics, Osaka University, Osaka 567-0047, Japan

Transitions among excited nucleons are studied within a non-relativistic quark model with a deformed harmonic oscillator potential. The transition amplitudes are factorized into the l -th moment and a geometrical factor. This fact leads to an analogous result to the “Alaga-rule” for baryons.

1. Introduction

There are many phenomenological models for masses and/or transition properties of baryons. However, if one tries to extend a model from the flavor SU(2) to SU(3) or to include excited state, the situation of fitting the experimental data becomes more and more complicated. Even if the spectrum can be reproduced by a phenomenological model, it is not easy to extract underlying physics or intuitive picture from them since some of the important effects may be hidden behind the model parameters. This is due to the seemingly complicated structure of the experimental data. It seems important therefore to have a fresh look at the existing data.

Historically, it is known that the Gell-Mann-Okubo (GO) mass formula works well to describe the pattern of the flavor octet and decuplet baryons in the ground states. We have extended this GO mass formula to all observed flavor SU(3) baryons and found that there is a flavor independent systematics in spectra [2, 3], more precisely in the excitation energy. Remarkably, we have found that these patterns of mass spectra have a quite similar behavior to the *rotational band* in the deformed nuclei, which seem to imply spatial deformation of excited states of baryons. We have then investigated these spectra in terms of an effective non-relativistic quark model with a deformed harmonic oscillator potential, which we call the deformed oscillator quark (DOQ) model, and have shown that almost all data can be reproduced with only one parameter by the DOQ model.

In the deformed nuclei, typical transitions are observed among rotational bands, which are classified as *inter*-band or *intra*-band transitions. We expect that we can also observe similar transitions in the excited baryons if the pattern of their spectra is described by rotational bands. In this brief report, we study intra-band transitions which correspond to the transitions among excited baryons with the same parity through an emission of a pion in the framework of the DOQ model.

*Presented by M. Koma at “PaNic02”, Osaka, Japan, 30 Sep. - 4 Oct. 2002

2. The DOQ model

Let us start with the hamiltonian of the DOQ model:

$$H_{\text{DOQ}} = \sum_{i=1}^3 \left(\frac{\vec{p}_i^2}{2m} + \frac{1}{2}m(\omega_x^2 x_i^2 + \omega_y^2 y_i^2 + \omega_z^2 z_i^2) \right), \quad (1)$$

where $\vec{p}_i$ and $\vec{r}_i = (x_i, y_i, z_i)$ are the momentum and position of the i -th quark. The constituent quark mass m is taken to be 300 MeV, which is insensitive to the following calculations. In order to take into account the deformation of excited states, the oscillator parameters $\omega_{x,y,z}$ can take different values, which are determined by energy minimization $\partial E/\partial\omega_x = 0, \partial E/\partial\omega_z = 0$, with volume conservation condition $\omega_x\omega_y\omega_z \equiv \omega^3 = (\text{const.})$, where E is the eigenenergy of H_{DOQ} . The only one parameter to be fitted in the DOQ model is ω , which determines the energy scale of the system. We choose $\omega = 644$ MeV to reproduce the excitation energy of the first $1/2^+$ excited states. The system is characterized by the total principle quantum number $N = n_x + n_y + n_z$, where n 's are oscillator quanta along each axis. Here we take $N = n_z, n_x = n_y = 0$, since this combination gives the lowest energy for given N . Resulting system is spherical for $N = 0$ (ground state) and prolately deformed: 1:2 for $N = 1$ (excited states with negative parity) 1:3 for $N = 2$ (those with positive parity).

Inspired by such a large deformation of the system, we adopt semi-classical picture to obtain the orbital wave function with a definite angular momentum L and its z -component L_z ; we introduce collective motion of strongly deformed intrinsic state by Wigner D function. Together with the appropriately symmetrized spin and isospin wave function χ and ϕ , the DOQ wave function with total spin J and its z -component M is written as $\Psi^{N;JM}(\vec{r}) = \psi^{JM}(\vec{\Theta})\varphi_{\text{int}}^N(\vec{r}_{\text{int}})$, where $\psi^{JM} \equiv [D_{0L_z}^L(\vec{\Theta})\chi]_{JM}$. Here the collective coordinate and internal body fixed coordinate are denoted as $\vec{\Theta}$ and $\vec{r}_{\text{int}}$.

3. Intra-band transitions

We calculate transition amplitudes between two excited nucleons (N^*) with a pion, $T_{N^*(JM) \rightarrow \pi N^*(J'M')}$. For $H_{\pi qq}$, we take the standard non-relativistic πqq interaction,

$$H_{\pi qq}(\vec{r}) = -\frac{g}{2m} \frac{1}{\sqrt{2\omega_k}} \left(\vec{\sigma} \cdot \vec{\nabla} \exp(i\vec{k} \cdot \vec{r}) \right) \tau \quad (2)$$

where k and ω_k are the momentum and energy of an emitted pion, respectively. Depending on the charge of an emitted pion, an appropriate component of the Pauli matrix for isospin τ acts on the isospin wave function of the N^* state.

Here we adopt a semi-classical cranking method to calculate the transition amplitudes;


$$T_{N^*(JM) \rightarrow \pi N^*(J'M')} = \int d^3\Theta \psi^{*J'M'}(\vec{\Theta}) \mathcal{O}_{\text{int}}(\vec{\Theta}) \psi^{JM}(\vec{\Theta}), \quad (3)$$

$$\mathcal{O}_{\text{int}}(\vec{\Theta}) = \int [d^3r_{\text{int}}] \varphi_{\text{int}}^{*N}(\vec{r}_{\text{int}}) R^\dagger(\vec{\Theta}) H_{\pi qq}(\vec{r}) R(\vec{\Theta}) \psi_{\text{int}}^N(\vec{r}_{\text{int}}), \quad (4)$$

where $R(\vec{\Theta})$ is a rotation operator which relates the original coordinate $\vec{r}$ to the internal body-fixed coordinate $\vec{r}_{\text{int}}$. Resulting transition amplitudes are written in terms of the l -th moment $Q^{(l)}(k) \equiv \int [d^3r_{\text{int}}] \varphi_{\text{int}}^{*N}(\vec{r}_{\text{int}}) j_l(kr) Y_{l0}(\hat{r}) \varphi_{\text{int}}^N(\vec{r}_{\text{int}})$, and some geometrical factors.

Table 1

Selection rules for strong intra-band decays $N^* \rightarrow N^*\pi$ (upper: positive parity, lower: negative parity)



L	J^P	J'^P	l_π	κ	κ'
$L = 4$	$9/2^+ \rightarrow$	$5/2^+$	3	$+$	$+$
	$9/2^+ \rightarrow$	$3/2^+$	3	$+$	$-$
	$7/2^+ \rightarrow$	$5/2^+$	1, 3	$-$	$+$
	$7/2^+ \rightarrow$	$3/2^+$	3^+	$-$	$-$
$L = 2$	$5/2^+ \rightarrow$	$1/2^+$	3	$+$	$+$
	$3/2^+ \rightarrow$	$1/2^+$	1, 3	$-$	$+$
L	J^P	J'^P	l_π	κ	κ'
$L = 3$	$7/2^- \rightarrow$	$3/2^-$	3	$+$	$+$
	$7/2^- \rightarrow$	$1/2^-$	3	$+$	$-$
	$5/2^- \rightarrow$	$3/2^-$	1, 3	$-$	$+$
	$5/2^- \rightarrow$	$1/2^-$	3	$-$	$-$

The definition of the decay width W is

$$W = 6\pi \int \frac{d^3k}{(2\pi)^3} \delta(E' + \omega_k - E) \frac{1}{2J+1} \sum_{MM'} |T_{N^*(JM) \rightarrow \pi N^*(J'M')}|^2, \quad (5)$$

where $E(E')$ is the energy of the initial (final) N^* state. Since both initial and final states are resonances, 1) average of the initial spin states, 2) summation over the final spin states, 3) integration over the momentum of emitted pions and 4) an overall factor three for the final state of pions (π^+ , π^- , π^0) are taken into account.

Since the difference between angular momentum of initial and final states is always two for the intra-band transitions, we can label transitions by L , l_π and relative orientation of the spin and the orbital angular momentum for the initial (final) state, κ (κ'). At the same time, the conservation of the angular momentum and p -wave coupling of pion lead to the selection rules as shown in Table 1.

After some straightforward calculations, the decay width (5) can be written as

$$W = \frac{27g^2}{2m^2} \tilde{k} \sum_{l_\pi} \left| F_{l_\pi; L}^{(N)\kappa\kappa'}(\tilde{k}) \right|^2, \quad \tilde{k} = \sqrt{(E_i - E_f)^2 - m_\pi^2}, \quad (6)$$

where we define transition form factor $F_{l_\pi; L}^{(N)\kappa\kappa'}$, where $\kappa(\kappa')$ is either $+$ or $-$.

As an example, we present a transition form factor for $L - 1/2 \rightarrow L' + 1/2$ transitions, which is dominated by $l_\pi = 1$ pion:

$$F_{l_\pi=3; L}^{(N)-+}(k) + F_{l_\pi=1; L}^{(N)-+}(k) \sim F_{l_\pi=1; L}^{(N)-+}(k) = i^1 (-1)^L k \frac{\sqrt{L-1}}{\sqrt{2L-1}} \left[\frac{1}{\sqrt{5}} Q_N^{(2)}(k) \right]. \quad (7)$$

From this expression, an interesting relation among transitions is obtained; apart from the effect of momentum transfer $\tilde{k}$, the ratio of the decay widths is determined only by

Table 2

Decay widths for each transition are shown in units of MeV. The classification scheme of the column is based on the selection rules given in Table 1 .

$L = 4 \rightarrow 2$	$9/2^+ \rightarrow 5/2^+$	$9/2^+ \rightarrow 3/2^+$	$7/2^+ \rightarrow 5/2^+$	$7/2^+ \rightarrow 3/2^+$
	5.8	10.8	35.3	1.7
$L = 2 \rightarrow 0$	$5/2^+ \rightarrow 1/2^+$	-	$3/2^+ \rightarrow 1/2^+$	-
	0.1	-	0.2	-
$L = 3 \rightarrow 1$	$7/2^- \rightarrow 3/2^-$	$7/2^- \rightarrow 1/2^-$	$5/2^- \rightarrow 3/2^-$	$5/2^- \rightarrow 1/2^-$
	7.9	9.9	39.3	0.7

geometrical factors in Eq. (7) as

$$\frac{W(L+2-1/2 \rightarrow L+1/2)}{W(L-1/2 \rightarrow L-2+1/2)} = \frac{(L+1)(2L-1)}{(2L+3)(L-1)}, \quad \frac{W(7/2 \rightarrow 5/2)}{W(3/2 \rightarrow 1/2)} = \frac{9}{7}. \quad (8)$$

This relation corresponds to the “Alaga rule” in nuclear physics [4].

Numerical results of the decay width for other transitions are listed in Table2.

4. Summary and discussion

We have found several interesting features among the intra-band transitions: 1) the decay widths of $L=2$ to $L=0$ are very small due to the small momentum transfer and hence a small phase space; we can show W has large k dependence, roughly $W \propto k^7$, with the long wave length approximation ($kr \ll 1$). 2) the decay widths of $7/2^+ \rightarrow 5/2^+$ and of $5/2^- \rightarrow 3/2^-$ have a rather large value since the pion with $l_\pi=1$ can contribute them (See, Table 1). Due to the difference of the geometrical factor in the transition form factor F the contribution of the $l_\pi=1$ transition is much bigger than that of $l_\pi=3$ transition. 3) the decay widths of $7/2^+ \rightarrow 3/2^+$ and of $5/2^- \rightarrow 1/2^-$ are small again due to the geometrical factor in the transition form factor. We would like to mention that such a interpretation of the result is possible by virtue of the semi-classical cranking formula.

Experimentally, such intra-band transitions may be observed in $N^* \rightarrow N\pi\pi$ processes. However, no clear evidence has been found in existing data at the moment. As a future work, we would like to study other transition properties such as inter-band transitions with a pion, which can be comparable with experimental data.

REFERENCES

1. D.E. Groom *et al.* (Particle Data Group), Eur. Phys. J. C15 (1 - 4) (2000) 1.
2. M. Takayama, H. Toki, A. Hosaka, Prog. Theor. Phys. 101 (1999) 1271.
3. M. Takayama, A. Hosaka, H. Toki, Nucl. Phys. A663&664 (2000) 695c.
4. G. Alaga and V. Paar, Phys. Lett. 61B (1976) 129.